

**New/Changed Dose**

Prescriber Information											
Prescriber Name:					MD	DO	NP	PA	NPI:		
Office Contact:					Practice Name / Collaborating MD:						
Address:				City:			State:		Zip:		
Phone:			Fax:								
Patient Information • PLEASE SEND COPY OF INSURANCE CARD											
Patients Name:			Last 4 Digits of SS#:		DOB: / /		Sex: M F		Weight: Height: Diabetic? Y N		
Address:				City:			State:		Zip:		
Home Phone:			Work/Cell:		HIPAA Contact:			Emergency #:			
Interpreter Needed? Y N		Allergies: Y N <b>If Yes, list allergies:</b>									
Insurance Information											
Primary Insurance:			Policy ID:		Group #:		BIN:		PCN:		
Policyholder Name:					Policyholder DOB: / /						
Dermatology • PLEASE SEND COPY OF MEDICAL AND PRESCRIPTION INSURANCE CARDS, PROGRESS NOTES AND LAB REPORTS, SUPPORTING DIAGNOSIS, CO-MORBIDITIES AND LAB VALUES											
ICD-10/Diagnosis Code:		Psoriasis Vulgaris (L40.0)		Other Psoriasis (L40.8)		Psoriasis unspecified (L40.9)		Psoriatic Arthritis (L40.5)		Hidradenitis Suppurativa (L73.2) Chronic Urticaria (L50.8)	
Atopic Dermatitis (L20.9)		Basal cell carcinoma (C44. _____)		TB/PDD Test Given: Y N		Date of Neg. Test: / /		HBV Positive? Y N		<b>If Yes, is patent currently treated? Y N</b>	
Prior Treatment? Y N (Provide Information Below)		BSA Affected (%):		Affected Areas: Palms Soles Head Neck Genitalia Other:							
Gastroenterology • PLEASE SEND COPY OF MEDICAL AND PRESCRIPTION INSURANCE CARDS, PROGRESS NOTES AND LAB REPORTS, SUPPORTING DIAGNOSIS, CO-MORBIDITIES AND LAB VALUES											
ICD-10/Diagnosis Code:		Crohn's Disease: K50.0 (Crohn's of the Small Intestine)		K50.1 (Crohn's of the Large Intestine)		K50.8 (Crohn's of Both Intestines)		K50.9 (Crohn's, Unspecified)			
Ulcerative Colitis: K51.0 (Ulcerative Pancolitis)		K51.2 (Ulcerative Procolitis)		K51.3 (Ulcerative Rectosigmoiditis)		K51.5 (Left Sided Colitis)		K51.8 (Other Ulcerative Colitis)			
K51.9 (Ulcerative Colitis, Unspecified)		K58.0 (Irritable Bowel Syndrome with Diarrhea)		Other:							
Date of Diagnosis: / /				Date of Negative TB Test: / /			Prior Treatment? Y N (Provide Information Below)				
Rheumatology • PLEASE SEND COPY OF MEDICAL AND PRESCRIPTION INSURANCE CARDS, PROGRESS NOTES AND LAB REPORTS, SUPPORTING DIAGNOSIS, CO-MORBIDITIES AND LAB VALUES											
Diagnosis:		M32.9 Active Systemic Lupus Erythematosus		M45.9 Ankylosing Spondylitis		M08.0 Juvenile Idiopathic Arthritis		L40.59 Psoriatic Arthritis		L40.54 Psoriatic Juvenile Arthritis	
M06.9 Rheumatoid Arthritis		M45.A _____ Non-Radiographic Axial Spondyloarthritis		Other:							
Date Diagnosis: / /		Date of Neg. TB Test: / /		Any prior treatment? Y N <b>If Yes, provide information below:</b>							
Prior Therapy											
Prior Therapy:					Reason for Discontinuation of Therapy:			Approx. Start Date: / /		Approx. End Date: / /	
Comorbidities:				Concomitant Medications:			Allergies: NKDA Other:				
Prescription Information											
Medication	Quantity/Dose		Sig					Refills			
<b>REMICADE®</b>	Number of 100mg vials: _____		<b>Starter Dose:</b> Infuse _____mg IV over 2 hours at weeks 0, 2 and 6					<b>No Refills</b>			
			<b>Maintenance Dose:</b> Infuse _____mg IV over 2 hours once every _____ weeks								
<b>RITUXAN®</b>	Number of 100mg/10ml vials: _____		<b>Starter Dose:</b> Infuse 1000mg IV over 4-6 hours on day 1 and day 15					<b>No Refills</b>			
	Number of 500mg/50ml vials: _____		<b>Maintenance Dose:</b> Infuse 1000mg IV over 4-6 hours every _____ weeks								
<b>SIMPONI ARIA®</b>	Number of 50mg/4ml vials: _____		<b>Starter Dose:</b> Infuse _____ mg IV over 30 minutes at weeks 0 and 4					<b>No Refills</b>			
			<b>Maintenance Dose:</b> Infuse _____ mg IV over 30 minutes once every 8 weeks								
<b>SKYRIZI®</b>	1 Vial (600mg/10ml)		<b>Starter Dose:</b> Infuse 600mg IV over at least 1 hour at week 0, week 4 and week 8. Begin SQ maintenance regimen at week 12					<b>2 Refills</b>			
<b>STELARA®</b>	Number of 45mg/0.5ml vials: _____		<b>Starter Dose:</b> Begin the SQ maintenance regimen 8 weeks after the initial IV dose					<b>No Refills</b>			
	Number of 130mg/26ml vials: _____		Weight > 85kg: Infuse 520mg IV over 1 hour Weight 56kg - 85kg: Infuse 390mg IV over 1 hour Weight ≤ 55kg: Infuse 260mg IV over 1 hour								

By signing this form and utilizing our services, you are authorizing Meijer and its employees to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.

Prescriber Signature	Date	Prescriber Signature	Date
----------------------	------	----------------------	------

Substitution Permitted

Dispense as Written

If brand is required, please write "DAW" in the box to the right.