

general nutrition recommendations

Psoriatic Disease

A balanced diet is important for everyone, but with psoriatic disease (PD) certain food choices may worsen symptoms. No one diet is proven best for those with PD, but below are tips for better managing the condition.

Nutrition Recommendations	
3 Meals a Day Aim to nourish your body daily. A minimum of three meals every day is recommended to keep energy levels high.	Eat Snacks Snacking between meals may help bridge the hunger gap if meals are eaten more than four hours apart.
Eat Small Meals If experiencing stomach ache or discomfort, it may be easier to eat several smaller meals throughout the day.	Colorful Foods Eat a variety of different colors and types of fruits and vegetables every day.
Whole Grains Include a variety of whole grain products every day: brown rice, oats, whole grain bread, whole grain pasta, quinoa, barley, and rye.	Lean Proteins Include a lean protein at every meal: chicken, fish, low-fat dairy products, beans, or tofu.
Include Healthy Fats Include a variety of food sources high in omega-3 fatty acids and monounsaturated fats: almonds, walnuts, olives, olive oil, canola oil, salmon, albacore tuna, chia seeds, flax seeds, and avocado.	Healthy Probiotics Aim to add food sources of probiotics in the diet for a wide range of useful bacteria. Examples include kombucha, sauerkraut, kimchi, kefir, coconut kefir, and yogurt.
Limit Unhealthy Fats Limit food sources high in saturated and trans fats: pastries, doughnuts, cakes, cookies, red meat, and full-fat dairy products such as ice cream.	Stay Hydrated Hydrate your body with water. Limit intake of soft drinks, sweetened teas, caffeinated beverages, energy drinks, and juice.

COMMON PROBLEMATIC FOODS

Common food allergens and sensitivities

Eggs, dairy, peanuts, nuts, soy, fish, shellfish, wheat, and gluten.

Nightshades

A class of vegetables, including tomatoes, peppers, eggplant, and potatoes.

FODMAPs

Naturally occurring sugars that are hard for your body to break down.

Gluten

Primarily found in foods made with wheat, rye, and barley.

Refined sugars

Cane juice, corn syrup, high-fructose corn syrup, raw sugar, crystal solids, brown rice syrup, maple syrup, honey, and brown sugar

Sugar alcohols and artificial sweeteners

Sorbitol, mannitol, xylitol, malitol - these are often found in products labeled "sugar free", "no sugar added", or "diet".

Processed and smoked meats

Hot dogs, bratwursts, sausages, highly-processed deli meats, and jerky.

Food additives

(found on ingredient lists of packaged products)
Carrageenan and maltodextrin

*Problematic foods are different for each patient but have been known to increase swelling or cause joint pain when eaten on a regular basis.