

Prescriber Information							
Prescriber Name:			MD	DO	NP	PA	NPI:
Office Contact:				Practice Name / Collaborating MD:			
Address:		City:		State:		Zip:	
Phone:		Fax:					

Patient Information • PLEASE SEND COPY OF INSURANCE CARD								
Patients Name:		Last 4 Digits of SS#:		DOB: / /	Sex: M F	Weight:	Height:	Diabetic? Y N
Address:			City:		State:		Zip:	
Home Phone:		Work/Cell:		HIPAA Contact:		Emergency #:		
Interpreter Needed? Y N	Allergies: Y N If Yes, list allergies:							

Insurance Information					
Primary Insurance:		Policy ID:	Group #:	BIN:	PCN:
Policyholder Name:			Policyholder DOB: / /		

Clinical Information • PLEASE SEND COPY OF MEDICAL AND PRESCRIPTION INSURANCE CARDS, PROGRESS NOTES AND LAB REPORTS, SUPPORTING DIAGNOSIS, CO-MORBIDITIES AND LAB VALUES							
ICD-10 Code:		Weight: lb / kg	Height: in / cm	BSA	m2	Diagnosis Date: / /	
Current Scr	or current GFR	ml/min	Confirmed Mutations:				

Notes to Pharmacy

Prescription Information				
Medication	Dose/Strength	Sig	Quantity	Refills
AFINITOR® (everolimus)	2.5mg tablet 5mg tablet	7.5mg tablet 10mg tablet		
AFINITOR DISPERZ® (everolimus for oral suspension)	2mg tablet 3mg tablet 5mg tablet			
CYCLOPHOSPHAMIDE	25mg capsule 50mg capsule 50mg tablet			
FASLODEX® (fulvestrant)	250mg/5ml PFS (2 per carton)	Starter Dose: Inject 500mg IM into the buttocks on days 1 and 15. Begin maintenance dose on day 29. Other:	2 Cartons	No Refills
	250mg/5ml PFS (2 per carton)	Maintenance Dose: Inject 500mg IM into the buttocks once monthly. Other:		
HERCEPTIN HYLECTA™ (trastuzumab and hyaluronidase-oysk)	5ml single dose vial (120mg trastuzumab, 2,000u hyaluronidase/ml)	Inject 5ml (600mg/10,000u) SQ once every 3 weeks Other:		
INLURIYO™ (imlunestrant)	200mg tablet	Take 2 tablets (400mg) by mouth daily on an empty stomach Other:		
KEYTRUDA QLEX™ (pembrolizumab and berahyaluronidase alfa-pmph)	2.4ml single dose vial (165mg pembrolizumab, 2,000u berahyaluronidase alfa/ml) 4.8ml single dose vial (165mg pembrolizumab, 2,000u berahyaluronidase alfa/ml)	Inject 2.4ml (395mg/4,800u) SQ once every 3 weeks Inject 4.8ml (790mg/9,600u) SQ once every 6 weeks Other:		
PHESGO™ (pertuzumab, trastuzumab and hyaluronidase-zzxf)	15ml single dose vial (80mg, 40mg, and 2,000u/ml)	Starter Dose: Inject 15ml SQ Other:		No Refills
	10ml single dose vial (60mg, 60mg, and 2,000u/ml)	Maintenance Dose: Inject 10ml SQ every 3 weeks Other:		
TYKERB® (lapatinib)	250mg tablet	Take 1,500mg (6 tablets) by mouth once daily without food (at least 1 hour before or 1 hour after a meal) Take 1,250mg (5 tablets) by mouth once daily without food (at least 1 hour before or 1 hour after a meal) Other:		
XELODA® (capecitabine)	150mg tablet 500mg tablet			

By signing this form and utilizing our services, you are authorizing Meijer and its employees to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.

Prescriber Signature	Date	Prescriber Signature	Date
----------------------	------	----------------------	------

Substitution Permitted

Dispense as Written

If brand is required, please write "DAW" in the box to the right.