

| Prescriber Information |  |      |       |                                   |        |    |      |      |
|------------------------|--|------|-------|-----------------------------------|--------|----|------|------|
| Prescriber Name:       |  |      |       | MD                                | DO     | NP | PA   | NPI: |
| Office Contact:        |  |      |       | Practice Name / Collaborating MD: |        |    |      |      |
| Address:               |  |      | City: |                                   | State: |    | Zip: |      |
| Phone:                 |  | Fax: |       |                                   |        |    |      |      |

| Patient Information • PLEASE SEND COPY OF INSURANCE CARD |  |                       |       |                |          |              |         |               |
|--|--|-----------------------|-------|----------------|----------|--------------|---------|---------------|
| Patients Name:   |  | Last 4 Digits of SS#: |       | DOB: / /       | Sex: M F | Weight:      | Height: | Diabetic? Y N |
| Address:   |  |                       | City: |                | State:   |              | Zip:    |               |
| Home Phone:  |  | Work/Cell:            |       | HIPAA Contact: |          | Emergency #: |         |               |
| Interpreter Needed? Y N                                  | Allergies: Y N If Yes, list allergies: |                       |       |                |          |              |         |               |

| Insurance Information |  |            |  |                       |  |      |  |      |
|-----------------------|--|------------|--|-----------------------|--|------|--|------|
| Primary Insurance:    |  | Policy ID: |  | Group #:              |  | BIN: |  | PCN: |
| Policyholder Name:    |  |            |  | Policyholder DOB: / / |  |      |  |      |

| Clinical Information • PLEASE SEND COPY OF MEDICAL AND PRESCRIPTION INSURANCE CARDS, PROGRESS NOTES AND LAB REPORTS, SUPPORTING DIAGNOSIS, CO-MORBIDITIES AND LAB VALUES                  |  |  |                               |  |  |                         |  |
|---|--|--|-------------------------------|--|--|-------------------------|--|
| ICD-10/Diagnosis Code:  |  | Crohn's Disease: K50.0__ (Crohn's of the Small Intestine) K50.1__ (Crohn's of the Large Intestine) K50.8__ (Crohn's of Both Intestines) K50.9__ (Crohn's, Unspecified) |                               |  |  |                         |  |
| Ulcerative Colitis: K51.0__ (Ulcerative Pancolitis) K51.2__ (Ulcerative Procolitis) K51.3__ (Ulcerative Rectosigmoiditis) K51.5__ (Left Sided Colitis) K51.8__ (Other Ulcerative Colitis) |  |  |                               |  |  |                         |  |
| K51.9__ (Ulcerative Colitis, Unspecified) K58.0__ (Irritable Bowel Syndrome with Diarrhea) Other:   |  |  |                               |  |  |                         |  |
| Date of Diagnosis: / /  |  |  | Date of Negative TB Test: / / |  | Prior Treatment? Y N (Provide Information Below) |                         |  |
| Prior Therapy:  |  | Reason for Discontinuation of Therapy:   |                               |  |  | Approx. Start Date: / / |  |
|   |  |  |                               |  |  | Approx. End Date: / /   |  |

| Prescription Information                          |  |  |                   |
|---|--|--|-------------------|
| Medication  | Quantity/Dose  | Sig  | Refills           |
| <b>SIMPONI®</b><br>SmartJect PFS                  | 3 cartons (100mg/ml)   | <b>Starter Dose:</b> Inject 200mg SQ at week 0; then 100mg at week 2   | <b>No Refills</b> |
|   | 1 carton (100mg/ml)  | <b>Maintenance Dose:</b> Inject 100mg SQ every 4 weeks, starting at week 6   |                   |
| <b>SKYRIZI®</b>                                   | 1 cartridge (360mg/2.4ml) with on-body injector<br>1 cartridge (180mg/1.2ml) with on-body injector | <b>Maintenance Dose:</b><br>Inject 360mg SQ beginning at week 12, and every 8 weeks thereafter<br>Inject 180mg SQ beginning at week 12, and every 8 weeks thereafter |                   |
| <b>STELARA®</b>                                   | 1 carton (1x90mg/ml PFS)   | <b>Maintenance Dose:</b> Inject 1ml (90mg) SQ 8 weeks after infusion, then every 8 weeks thereafter  |                   |
| <b>TREMFYA®</b><br>PFS Pen                        | CD Induction Pack (2x200mg/ml PENS)  | <b>CD Starter Dose:</b> Inject 400mg SQ at weeks 0, 4 and 8  | <b>2 Refills</b>  |
|   | 1 carton (1x100mg/ml)<br>1 carton (1x200mg/ml)   | <b>Maintenance Dose:</b><br>Inject 100mg SQ at week 16 and every 8 weeks thereafter<br>Inject 200mg SQ at week 12 and every 4 weeks thereafter                       |                   |
| <b>VELSIPITY™</b>                                 | 2mg tablets (30 day supply)  | Take 1 tablet by mouth once daily  |                   |
| <b>XELJANZ®</b>                                   | 10mg tablets (quantity QS for length of starter dose therapy, in multiples of 60 tablets)          | <b>Starter Dose:</b> Take 10mg by mouth twice daily for ____ weeks   | <b>No Refills</b> |
|   | 5mg tablets (30 day supply)<br>10mg tablets (30 day supply)  | <b>Maintenance Dose:</b> Take 1 tablet by mouth two times a day  |                   |
| <b>XELJANZ® XR</b>                                | 22mg tablets (quantity QS for length of starter dose therapy, in multiples of 30 tablets)          | <b>Starter Dose:</b> Take 22mg by mouth once daily for ____ weeks  | <b>No Refills</b> |
|   | 11mg tablets (30 day supply)<br>22mg tablets (30 day supply)                                       | <b>Maintenance Dose:</b> Take 1 tablet by mouth once daily   |                   |
| <b>XIFAXAN®</b>                                   | 200mg tablet<br>550mg tablet   | Take 1 tablet by mouth 2 times a day for ____ days<br>Take 1 tablet by mouth 3 times a day for ____ days   |                   |
| <b>ZEPOSIA®</b>                                   | Starter Pack (7 day supply)<br>Starter Kit (37 day supply)   | Take 0.23mg by mouth daily on days 1-4, then 0.46mg daily on days 5-7, then 0.92mg daily thereafter  |                   |
|   | 0.92mg capsules (30 day supply)  | <b>Maintenance Dose:</b> Take 1 capsule by mouth daily   |                   |
| <b>ZYMFENTRA™</b><br>(infliximab-dyyb)<br>PFS Pen | 1 carton (2x120mg/ml)  | <b>Maintenance Dose:</b> Inject 120mg SQ every 2 weeks, starting at week 10  |                   |

| Injection Training                  |   |   |
|-------------------------------------|---|---|
| Patient received injection training | Prescriber's office to provide injection training | Meijer to coordinate injection training |

By signing this form and utilizing our services, you are authorizing Meijer and its employees to serve as your prior authorization designated agent in dealing with medical and prescription insurance companies.

|                      |      |                      |      |
|----------------------|------|----------------------|------|
| Prescriber Signature | Date | Prescriber Signature | Date |
|----------------------|------|----------------------|------|

Substitution Permitted

Dispense as Written  
If brand is required, please write "DAW" in the box to the right.